arXiv:1409.8425v1 [hep-ph] 30 Sep 2014

High-Energy Scattering vs Static QCD Strings

V.A. Petrov[a1], R.A. Ryutin[b1]

Institute for High Energy Physics,*142 281*, Protvino, Russia

Abstract. We discuss the shape of the interaction region of the elastically scattered protons stipulated by the high-energy Pomeron exchange which turns out to be very similar with the shape of the static string representing the confining QCD flux tube. This similarity disappears when we enter the LHC energy region, which corresponds to many-Pomeron exchanges. Reversing the argument we conjecture a modified relationship between the width and the length of the confining string at very large lengths.

PACS. 11.55.Jy Regge formalism – 12.38.Aw General properties of QCD (dynamics, confinement, etc.) – 11.15.-q Gauge field theories – 12.40.Nn Regge theory, duality, absorptive/optical models – 11.25.Wx String and brane phenomenology – 11.10.Jj Asymptotic problems and properties – 11.15.Ha Lattice gauge theory

1 Itroduction

The basic element of the Regge framework to describe the high-energy hadron scattering is the well-known expression

$$A(s,t) = \beta(t)\left(\mathrm{e}^{-\frac{\mathrm{i}\pi}{2}}s\right)^{\alpha(t)} + \dots \qquad (1)$$

where $\alpha(t)$ is the leading (Pomeron) Regge trajectory. The average transverse (w. r. t. the collision axis) size of the interaction region is given by the formula

$$< \boldsymbol{b}^2 >= 4\alpha'(0)\ln(s) + \dots \qquad (2)$$

Thus, in the Regge-pole framework the transverse size of the interaction region widens with energy.

It is not difficult to obtain the average longitudinal range of interaction which appears to be

$$R = < z_1 - z_2 >=< \partial(\mathit{scattering\ phase})/\partial p_{\parallel}^{out} >= \\ = \pi < \alpha'(t) > \sqrt{s}/2 + \dots \qquad (3)$$

where $z_{1,2}$ are longitudinal coordinates of the scattered hadrons in the c.m.s., $\mathit{scattering\ phase} = -\pi\alpha(t)/2$, $t = -2p^2 + 2pp_{\parallel}^{out}$, $s = 4p^2 + 4m^2$.

In linear approximation for the trajectory, $\alpha(t) = \alpha(0) + \alpha'(0)t$, we obtain

$$R = \pi\alpha'(0)\sqrt{s}/2 + \dots \qquad (4)$$

The fast growth with energy of the longitudinal interaction range in diffractive scattering is known for a long time (see f.e. [1,2,3,4,5,7]).

We get the following relationship between transverse ($< \boldsymbol{b}^2 >$) and longitudinal (R) sizes of elastic hadron scattering (at large R)

$$< \boldsymbol{b}^2 >= 8\alpha'(0)\ln(R) + \dots \qquad (5)$$

So the transverse size of the interaction region increases logarithmically with the growth of the longitudinal interaction range. Pictorially at large R it looks as a prolate cigar-like region:



Fig. 1. Interaction region.

2 The Shape of Static Strings

In paper [6] the question of the thickness of a static string spanned by infinitely heavy quark-anti-quark pair was addressed. Trying different frameworks the authors came to the conclusion that (the square of)the size of the transverse quantum fluctuations of a string-like confining flux grows logarithmically with the string length R.

[a] *e-mail:* Vladimir.Petrov@cern.ch
[b] *e-mail:* Roman.Rioutine@cern.ch

This growth is thought to overcome the classical mechanism of the chromo-electric flux squeezing (dual Meissner effect).

Much later the same problem was studied in a more rigorous way. In particular, in Ref. [7,8] it was shown that in $d = 2 + 1$ dimensional space-time the confining QCD string connecting two static sources ($Q\bar{Q}$) has the square of half-width w^2 (in the middle of the string length), which grows with the length of the string R as

$$w^2 = \frac{d-2}{2\pi\sigma}\ln(R) + \dots \tag{6}$$

where σ is the string tension.

If we assume that at high energies the interaction region is close to axisymmetric form, then we have for its half-width squared

$$w^2_{scat} = \frac{< \boldsymbol{b}^2 >}{4} = 2\alpha'(0)\ln(R) + \dots \tag{7}$$

If to take into account the relation between the Regge slope and string tension

$$\alpha' = \frac{1}{2\pi\sigma} \tag{8}$$

then (6) and (7) coincide if $d = 3+1$, i.e. formally outside the scope of the derivation of (6) in Ref. [1], where $d = 2 + 1$.

However, in paper [9] the same problem was studied in 4-dimensional space. The authors of this paper used the same logarithmic relation with the coefficient B in front of $\ln(R)$ as one of two free parameters for the fitting the lattice data. Below we show their picture.



Fig. 2. The width squared of the $q\bar{q}$ confining string as function of its length [9].

3 Reggeon Exchanges and Strings

If we identify the "static" coefficient $B = 0.0762 \pm 0.0035$ fm^2 with $2\alpha'(0)$ then we obtain

$$\alpha'(0) = 0.9525 \pm 0.04375 \text{ GeV}^{-2} \tag{9}$$

This slope is not very far from the values of the secondary Regge trajectories which are usually associated with open $q\bar{q}$ strings. So if the high-energy scattering were defined by the meson trajectories we could conclude that the interaction region is being stretched with the energy growth as an open string.

We, however, know that actually it is not the meson (secondary Reggeon) but the Pomeron trajectory which dominates the high-energy behaviour. In this case the coefficient in front of $\ln(R)$ is four times smaller. How can it be explained?

Usually one argues that taking into account the closed string nature of the Pomeron we should get a twice smaller coefficient corresponding to $\alpha'_{\mathbb{P}} = 0.5$ GeV^{-2}. However the high-energy phenomenology supports in a very strong way even smaller slope:

$$\alpha'_{\mathbb{P}} \cong 0.25 \text{ GeV}^{-2} \tag{10}$$

If we assume that the closed (stretched) string shape follows Eq.(6) with 4 times smaller coefficient in front of $\ln(R)$ then we obtain the following estimate for the Pomeron slope:

$$\alpha'_{\mathbb{P}}(0) \simeq 0.24 \pm 0.01 \text{ GeV}^{-2}. \tag{11}$$

Actually the tension values of various kinds of closed strings in the lattice framework were considered in Ref. [10] with one of the results fairly close to that of high-energy phenomenology:

$$\alpha'_{\mathbb{P}}(lattice) \simeq 0.253(2) \text{ GeV}^{-2}. \tag{12}$$

This is suggestive for the conclusion that the interaction region of high-energy hadron scattering clearly bears stringy features similar to the static string configurations in spite of their physically extraordinary difference: a highly dynamic phenomenon against the static one. In the case of scattering we deal with fluctuating average length while in Eq. (6) it is fixed parameter.

4 Breakdown of Reggeon-String Similarity or String Shape Modification?

We have to notice that Regge-Pomeron formula (6) begins to dominate at energies not less than the ISR energies ($\sqrt{s} = 20 \div 50$ GeV), this corresponds to the minimal interaction length $R = 1.5$ fm, which is exactly at the end of the lattice data at Fig.2.

If we believe that the interaction region looks like a stretched string of non-zero transverse extent then we come to a striking conclusion that these extremely stretched configurations can survive without breaking up to monstruous lengths of order of 150 fm. This concerns high-mass ($\simeq 1$ TeV$/c^2$) stringy configurations.

According to the ratio σ_{el}/σ_{tot} measured by the TOTEM LHC Collaboration [11]

$$\sigma_{el}/\sigma_{tot}(\text{TOTEM, } 7 \text{ TeV}) = 0.257 \pm 0.005 \tag{13}$$

this means that the probability of the long string-like configuration survival (before breaking out into the two elastically scattered protons) is of order of 25%. Such a phenomenon would mean that confinement is fairly important not only at low-energy hadron physics but at high-energy scattering as well.

Our arguments are heavily based on Eq.(7). However it ceases to be adequate at very high energy though remains a basic building block (eikonal) for the full unitary amplitude in the Regge-eikonal framework. Latest experiments at the LHC show that the width w^2 as function of R significantly deviates from the simple logarithmic dependence in the LHC energy region (see Fig. 3 below).



Fig. 3. The length dependence of the half-width of the transverse interaction region of the elastic scattering of (anti)protons. $w^2_{scat} = \frac{<b^2>}{4} \approx 2\alpha'_{\mathbb{P}}(0)\ln(R) \approx (0.14\ \text{fm})^2 \ln(R)$. The red point is the measurement of the TOTEM Collaboration at 7 TeV. The data from the ALFA experiment (ATLAS) [12] gives a close value. (The figure is a modification of Fig.35 from Ref. [12])

At ultra-high energies the relationship between w^2 and R changes significantly as could be seen from the following asymptotic behavior:

$$< \boldsymbol{b}^2 > (ultra - high\ energy) \approx 4\left[\alpha_{\mathbb{P}}(0) - 1\right]\alpha'_{\mathbb{P}}(0)\left[\ln(s)\right]^2 + \ldots \quad (14)$$

Unfortunately we haven't yet the analytic expression for the phase of the scattering amplitude in this case and can't insist that the proportionality

$$R \sim \sqrt{s}$$

still holds. However to trace the trend we can use the asymptotic (actually at practically unavailable energies) expression for the scattering amplitude:

$$T(s,t) \approx \frac{4\pi \mathrm{i} s R_0(s)}{\sqrt{-t}} J_1(R_0(s)\sqrt{-t}), \quad (15)$$

$$R_0^2(s) = 4\Delta_{\mathbb{P}}\alpha'_{\mathbb{P}}(0)\left[\ln(s) - \mathrm{i}\pi/2\right]^2, \quad \Delta_{\mathbb{P}} = \alpha_{\mathbb{P}}(0) - 1. \quad (16)$$

We get for the interaction region scales:

$$R \approx \pi\alpha'_{\mathbb{P}}(0)\Delta_{\mathbb{P}}\sqrt{s}\ln(s) + \ldots, \quad (17)$$

$$< \boldsymbol{b}^2 > = 4\alpha'_{\mathbb{P}}(0)\Delta_{\mathbb{P}}\left[\ln(s)\right]^2 + \ldots. \quad (18)$$

If we try to assume that the similarity discussed above persist at higher energies (larger R) then we come to the following modification of the string width-length relationship:

$$w^2_{string} = 2\frac{\Delta_{\mathbb{P}}}{\pi\sigma_{\mathbb{P}}}\left[\ln(R)\right]^2 + \ldots, \quad (19)$$

where $\sigma_{\mathbb{P}}$ is the tension of the string corresponding to the Pomeron while parameter $\Delta_{\mathbb{P}}$ can be only explained when considering the trajectory deviation from the linear behavior at small unphysical values of the string mass (see e.g. Ref. [13]). We have also to note that the value of the tension $\sigma_{\mathbb{P}} = 1/(2\pi\alpha'_{\mathbb{P}}(0))$ may differ from the "quasi-classical" tension of the straight-line string at high string masses.



Fig. 4. Towards the similarity of multi-Reggeon exchanges and account of possible topologies of the string.

The asymptotic form of the scattering amplitude (15) is due to summation of infinitely many Reggeons. If to liken the Reggeon to a string (a closed one in case of the Pomeron) then this would correspond to summation in all possible genera (handles and holes) of the string spanned between the sources (see Fig. 4). We could conjecture that account of higher genera in the problem of the static string width dependent on its length leads to formula (19).

5 Conclusions

We have shown that the relationship of the transverse and longitudinal interaction ranges which follow from a one-Reggeon-exchange scattering amplitude is identical to that of the static confining string. Reversing the argument we conjecture on the basis of the Regge-eikonal asymptotic for the scattering amplitude that account of higher genera modify the dependence of the static string width on its length.

Aknowledgements

We would like to thank our colleagues A.A. Godizov, A.K. Likhoded and G.P. Pronko for discussions.

References

1. E.L. Feinberg, *Inelastic Diffractive Processes at High Energies* (In Russian), Soviet Physics: Uspekhi. UFN **LVIII,2**, 193 (1956)
2. V.N. Gribov, B.L. Ioffe and I.Ya. Pomeranchuk, *What is the range of interaction at high energies?*, Sov. J. Nucl. Phys. **2**, 549 (1966), Yad. Fiz. **2**, 768 (1965)
3. V.N. Gribov, *Space-Time Description of the Hadron Interaction at High Energy*, Sov. J. Nucl. Phys. **9**, 369 (1969); arXiv:hep-ph/0006158
4. F.E. Low, *A Model of the Bare Pomeron*, Phys. Rev. D **12**, 163 (1975)
5. B. Scrempp and F. Schrempp, *Strong Interactions - A Tunneling Phenomenon?*, Nucl. Phys. B **163**, 397 (1980)
6. M. Luscher, G. Munster and P. Weisz, *How Thick are Chromoelectric Flux Tubes?*, Nucl. Phys. B **180**, 1 (1981)
7. F. Gliozzi, M. Pepe, and U.-J. Wiese, *Width of the Confining String in Yang-Mills Theory*, Phys. Rev. Lett. **104**, 232001 (2010)
8. H.B. Meyer, *Transverse Structure of the QCD String*, Phys. Rev. D **82**, 106001 (2010)
9. N. Cardoso, M. Cardoso and P. Bicudo, *Inside the SU(3) quark-antiquark QCD flux tube: screening versus quantum widening*, Phys. Rev. D **88**, 054504 (2013)
10. H.B. Meyer and M.J. Teper, *Glueball Regge trajectories and the pomeron: A Lattice study*, Phys. Lett. B **605**, 344 (2005)
11. The TOTEM Collab., *Luminosity-independent measurements of total, elastic and inelastic cross-sections at* $\sqrt{s} =$ 7 *TeV*, CERN-PH-EP-2012-353, Europhys. Lett. **101**, 21004 (2013)
12. U. Amaldi, *Small-angle physics at the intersecting storage rings forty years later*, "40th Anniversary of the First Proton-Proton Collisions in the CERN Intersecting Storage Rings (ISR)"; U. Amaldi, P.J. Bryant, P. Darriulat and K. Hubner, CERN-2012-004 (CERN, Geneva, 2012), p.33
13. L.D. Soloviev, *Multimode relativistic string: Classical solutions and quantization*, Theor. Math. Phys. **136**, 1225 (2003); Teor. Mat. Fiz. **136**, 380 (2003)